



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q 1/68	A1	(11) International Publication Number: WO 00/58507 (43) International Publication Date: 5 October 2000 (05.10.00)
(21) International Application Number: PCT/GB00/01222 (22) International Filing Date: 30 March 2000 (30.03.00) (30) Priority Data: 9907344.7 30 March 1999 (30.03.99) GB 9919603.2 18 August 1999 (18.08.99) GB (71) Applicant (for all designated States except US): SOLEXA LTD. [GB/GB]; 38 Jermyn Street, London SW1Y 6DN (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): KLENERMAN, David [GB/GB]; University of Cambridge, Dept. of Chemistry, Lensfield Road, Cambridge CB2 1EW (GB). (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: POLYNUCLEOTIDE SEQUENCING (57) Abstract <p>A method is disclosed for determining the sequence of a polynucleotide. The sequence is determined by generating the complement of the polynucleotide using the polymerase reaction to extend a suitable primer, and characterising the successive incorporation of bases that generate the complement. The method requires the sequential addition of a composition comprising the different bases A, T, G and C, a minor proportion of which are detectably-labelled.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

POLYNUCLEOTIDE SEQUENCING

Field of the Invention

This invention relates to the sequencing of polynucleotides. In particular, this invention discloses methods for determining the sequence of arrayed polynucleotides.

5 Background to the Invention

Advances in the study of molecules have been led, in part, by improvement in technologies used to characterise the molecules or their biological reactions. In particular, the study of the nucleic acids DNA and RNA has benefited from developing technologies used for sequence analysis and the study of hybridisation events.

10 An example of the technologies that have improved the study of nucleic acids, is the development of fabricated arrays of immobilised nucleic acids. These arrays consist typically of a high-density matrix of polynucleotides immobilised onto a solid support material. Fodor *et al*, Trends in Biotechnology (1994) 12:19-26, describes ways of assembling the nucleic acids using a chemically sensitized glass surface
15 protected by a mask, but exposed at defined areas to allow attachment of suitably modified nucleotide phosphoramidites. Fabricated arrays may also be manufactured by the technique of "spotting" known polynucleotides onto a solid support at predetermined positions (e.g. Stimpson *et al* PNAS (1995) 92:6379-6383).

A further development in array technology is the attachment of the
20 polynucleotides to the solid support material via beads (microspheres).

For DNA arrays to be useful their sequences must be determined. US 5302509 discloses a method to sequence polynucleotides immobilised on a solid support. The method relies on the incorporation of 3'-blocked bases A, G, C and T having a different fluorescent label to the immobilised polynucleotide, in the presence
25 of DNA polymerase. The polymerase incorporates a base complementary to the target polynucleotide, but is prevented from further addition by the 3'-blocking group. The label of the incorporated base can then be determined and the blocking group removed to allow further polymerisation to occur.

However, the need to remove the blocking groups after each cycle is time-
30 consuming and must be performed with high efficiency.

Similarly, EP0640146 discloses a polymerisation-based technique for sequencing DNA. The technique again requires removal of a blocking group prior to subsequent incorporation of nucleotides.

There is therefore a need for alternative methods for determining the sequence of arrayed polynucleotides.

Summary of the Invention

In the general method of the invention, a target polynucleotide sequence can
5 be determined by generating its complement using the polymerase reaction by the
extension of a suitable primer, and characterising the successive incorporation of
bases that generate the complement. The method requires the target sequence to be
immobilised on a solid support, with multiple copies of the target being localised within
discrete regions. Each of the different bases A, T, G or C are then brought, by
10 sequential addition, into contact with the target, and any incorporation events
detected. Repeating the procedure with each of the bases allows the sequence of the
complement to be identified, and thereby the target sequence also.

A distinguishing feature from the disclosure in US 5302509 is that the bases
do not contain a blocking group preventing further polymerisation from occurring. In
15 addition, the present invention requires the separate and serial addition of each of the
different base types to the array, and, when fluorophores are used as the label,
removal of the label can be carried out efficiently by photobleaching.

A further distinguishing feature, particularly relevant to EP 0640146, is that for
each incorporation step, only a minor proportion of the bases are detectably-labelled.
20 Consequently, among the many copies of the target, relatively few will incorporate a
labelled base into the complement. This permits the straight forward identification of
any sequence containing two or more consecutive bases of the same type. In this
case, copies of the target will incorporate differing amounts of the labelled base into
the complement, resulting in differing levels of signal. It is then possible to determine
25 quantitatively the number of consecutive bases on the complement by detecting the
different level of signals generated, as explained later.

Accordingly, a method for determining the sequence of a target polynucleotide
on an array, comprises the steps of:

- (i) forming an array comprising multiple copies of each target
30 polynucleotide;
- (ii) contacting the array with a composition comprising one of the bases
A, T, G or C under conditions that permit polymerisation to occur,
wherein a minor proportion of the bases are detectably-labelled;

- (iii) detecting the incorporation of a base onto the complement of the target after removal of non-incorporated bases; and
- (iv) repeating steps (ii) and (iii) with each of the different bases until the sequence is determined.

5 According to one aspect of the invention, the label from incorporated bases may be removed either prior to the addition of bases having the same label or before it becomes difficult to detect incorporation.

 According to a second aspect of the invention, when the label is a fluorophore, the fluorescence signal generated on nucleotide incorporation may be measured
10 quantitatively, without the need to remove labels after each incorporation step. There is therefore a method for determining the sequence of a target polynucleotide as described above, wherein the fluorescence labels are not removed from the incorporated nucleotides, and subsequent detection of incorporation is carried out by measuring the step wise increase in the fluorescence signal.

15 The advantage of this embodiment is that it does not require the step of photobleaching and may therefore be carried out quickly and efficiently.

 Sequencing the polynucleotides on the array makes it possible to form a spatially addressable array. This may then be used for many different applications, including genotyping studies and other characterisation experiments.

20 The method of the present invention may be automated to produce a very efficient and fast sequence determination.

Description of the Drawings

 Figure 1 represents a fluorescence (left) or optical (right) image generated in the presence (A) and absence (B) of polymerase enzyme; and

25 Figure 2 represents a fluorescence image generated from beads with fluorophore-labelled DNA attached (A) or a fluorophore-labelled nucleotide incorporated into DNA using a polymerase (B).

Description of the Invention

30 The method for determining the sequence of the arrayed polynucleotides is carried out by contacting the array separately with the different bases to form the complement to that of the target polynucleotide, and detecting incorporation. The method makes use of polymerisation, whereby a polymerase enzyme extends the complementary strand by incorporating the correct base complementary to that on the

target. The polymerisation reaction also requires a specific primer to initiate polymerisation.

For each cycle, adding one base type to the array, only a minor proportion of the bases are detectably-labelled, i.e. less than 50% of the bases are detectably-labelled, preferably less than 20%. Therefore, it is only the incorporation of detectably-labelled bases that can be monitored. The labelled bases are present at a fixed low concentration with respect to the non-labelled bases. The concentration may be chosen to permit a suitable incorporation rate of the labelled bases for efficient detection. For example the concentration may be chosen to permit between 10% to 0.0001% incorporation of labelled bases, preferably, between 5% and 0.01%, most preferably between 1% and 0.1%.

Using many copies of the same polynucleotide in discrete regions it is possible to detect quantitatively the incorporation of a labelled base. For example, on incorporation of the adenosine nucleotide, a proportion of the polynucleotides will have a non-labelled adenosine nucleotide and a proportion will have a labelled adenosine nucleotide. Detecting the incorporation of the label will allow a sequence determination to be made. If two adenosine nucleotides are incorporated consecutively into the complementary strand, a proportion of the polynucleotide copies will incorporate two non-labelled adenosine nucleotides, a proportion will incorporate one labelled adenosine and one non-labelled adenosine, and a proportion will incorporate two labelled adenosine nucleotides. However, the ratio of labelled to unlabelled nucleotide will be such that very little of the labelled nucleotide will incorporate into the same strand. This is especially preferable when fluorescent labels are used, where fluorescence quenching or loss of linearity of signal may be caused. The label will therefore be distributed throughout the population of a given sequence. Consequently, there will be a quantitative difference in the signal generated within the population of the given sequence. It is possible therefore to detect the incorporation of the two consecutive labelled bases due to the quantitative differences in the signal.

In the context of the invention, reference to the bases A, T, G and C is taken to be a reference to the deoxynucleoside triphosphates, Adenosine, Thymidine, Guanosine and Cytidine, and to functional analogs thereof, including dideoxynucleoside triphosphates.

The terms "arrayed polynucleotides" and "polynucleotide arrays" are used herein to define an array of polynucleotides that are immobilised on a solid support

material. The polynucleotides may be immobilised to the solid support indirectly through a linker molecule, or may be attached to a particle, e.g. a microsphere, which is itself attached to a solid support material.

5 An important requirement is that there are multiple copies of each target polynucleotide on the array. Typically, these will be in discrete positioned regions on the solid support. Each discrete region may typically comprise several hundred to several thousand copies of the target polynucleotide. There may be, for example, up to 10,000 polynucleotide copies per region. The polynucleotides within each region preferably form a substantially uniform arrangement. This permits a high level of
10 discrimination between individual polynucleotides, which may be preferable to resolve individual labels. However, it is not necessarily the density of the polynucleotides that is of primary importance; the concentration of the labelled bases during the sequencing steps is also important, and this can be optimised readily by the skilled person.

15 The term "spatially addressable" is used herein to describe how different molecules may be identified on the basis of their position on an array.

The detection of an incorporated base may be carried out by using a confocal scanning microscope to scan the surface of the array with a laser, to image a fluorophore bound directly to the incorporated base. Alternatively, a sensitive 2-D
20 detector, such as a charge-coupled detector (CCD), can be used to visualise the individual signals generated. The use of such apparatus is known to the skilled person. However, other techniques such as scanning near-field optical microscopy (SNOM) are available which are capable of smaller optical resolution, thereby committing "more dense" arrays to be used. For example, using SNOM, individual
25 polynucleotides may be distinguished when separated by a distance of less than 100 nm, e.g. 10 nm x 10 nm. For a description of scanning near-field optical microscopy, see Moyer *et al* Laser Focus World (1993) 29:10.

The polynucleotides that may be sequenced include DNA, RNA and synthetic alternatives such as PNA.

30 The polynucleotides may be attached to the solid support by recognised means, including the use of biotin-avidin interactions or the use of amine linkages. In one embodiment, the polynucleotides are attached to the solid support via microscopic beads (microspheres), which may in turn be attached to the solid support by known

means. The microspheres may be of any suitable size, typically in the range of from 10 nm to 100 nm in diameter.

Attachment via microspheres is a preferred embodiment as it allows discrete regions of polynucleotides to be easily generated on the array. Each microsphere
5 may have multiple copies of a polynucleotide attached, and each microsphere can be resolved individually to determine incorporation events.

The method makes use of the polymerisation reaction to generate the complementary sequence of the target. The conditions necessary for polymerisation to occur will be apparent to the skilled person. For example, a polymerase enzyme
10 may be used to extend the complementary strand, and different polymerases, including DNA polymerases and RNA polymerases, are known to those skilled in the art. For example, the Klenow fragment of E. coli DNA polymerase I or the T7 DNA polymerase may be used. To carry out the polymerase reaction it may be necessary to first anneal a primer sequence to the target polynucleotide, the primer sequence
15 being recognised by the polymerase enzyme and acting as an initiation site for the subsequent extension of the complementary strand. Other conditions necessary for carrying out the reaction, including temperature and pH, will be apparent to those skilled in the art.

This polymerisation step is allowed to proceed for a time sufficient to allow
20 incorporation of all the correct bases. This will depend on the efficiency of incorporation and can be determined by the skilled person. Bases that are not incorporated are then removed, for example, by subjecting the array to a washing step, and detection of the incorporated labels may then be carried out.

Detection may be by conventional means, for example if the label is a
25 fluorescent moiety, detection may be carried out by optical microscopy, e.g. confocal scanning microscopy.

A preferred embodiment of the invention uses fluorophores as the label, and many examples of fluorophores that may be used are known in the prior art e.g. tetramethylrhodamine (TMR).

30 After detection, the labels may be removed from the bases so that they do not interfere with the signal generated from next cycle of incorporation. If the label is a fluorophore it is possible to bleach the fluorophore by chemical means or through the use of a laser (photobleaching). Alternatively, the label may be removed by chemical or photochemical means.

The process of incorporating bases may then be repeated using each of the different bases until the sequence has been determined.

It may not always be necessary to remove the labels prior to the addition of the next base sample. Different bases may have distinguishable labels and so it will only
5 be necessary to remove incorporated labels prior to adding bases having an identical label.

In one embodiment, fluorescent labels are used and detection is carried out by optical means without the requirement for removing labels between incorporation steps.

10 For example, a confocal microscope may be used to scan the array and measure quantitatively the step-wise increase in fluorescence after each cycle of incorporation. By measuring the increase in the amount of fluorescence after each cycle, and not the absolute amount, it should be possible to determine whether there are two or more nucleotides incorporated consecutively onto the template. This
15 method relies on using sensitive detectors (e.g. charge coupled detectors) to measure the increase in signal. Suitable apparatus for carrying out the method is available commercially and will be apparent to the skilled person.

In a separate embodiment of the invention, the labelled bases may be modified so that on incorporation, no further bases may be added. Bases that carry out this
20 chain terminating function include the dideoxynucleoside triphosphates, as used in conventional Sanger sequencing (Proc. Natl. Acad. Sci. USA 74: 5463-5467, 1977).

Therefore, after each incorporation step, a proportion of the polynucleotides will incorporate a labelled base that prevents further chain-extension. The number of polynucleotides available for the polymerisation step will gradually decrease as the
25 sequencing method proceeds. However, provided there are sufficient copies of the polynucleotide, and provided the concentration of the labelled, chain-terminating bases, is sufficiently low, it should be possible to sequence the target polynucleotides.

The following experiment illustrates the invention.

Example

30 In this experiment a fluorescently-labelled DNA molecule (SEQ ID NO. 1) was coupled directly to beads and the level of fluorescence measured using an inverted Nikon microscope with an ICCD detector in an epifluorescence set-up. In a separate reaction, an unlabelled DNA (SEQ ID NO. 2) was attached to beads (containing SEQ ID NO. 2) and a fluorescently-labelled nucleotide incorporated onto the DNA (SEQ ID

NO. 2) using a polymerase. By comparing the average level of fluorescence between the two sets of beads the efficiency of incorporation of the fluorescently-labelled nucleotide was shown to be 89%. This was determined by diluting the fluorescent beads in unmodified beads so that each fluorescent bead could be detected individually.

By measuring the signal-to-noise in the experiment an estimate can be obtained of the fraction of nucleotides that can be labelled with a fluorophore and detected when incorporated. This is less than 1%, i.e. it is possible to detect incorporation when the concentration of fluorescently-labelled nucleotides is such that only 1% is incorporated, and the remaining 99% of the incorporated nucleotides are non-labelled.

The experiment is now described in more detail.

DNA Coupling

Carboxylic acid-modified beads (both non-porous polystyrene and silica) of sizes 0.5-2.9 μm were placed in solutions of milli-q water (typically 1 mg per 50 ml). 1-3(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1 mg) and the oligonucleotide (added to give a final concentration of 10 μM) were added, the beads agitated by vortexing and left for 12 h at room temperature. The beads were washed with 0.15 M NaOH, twice with TT buffer (250 mM Tris.HCl, pH 8.0, 0.1 % tween 20) and heated at 80°C in TTE (250 mM Tris.CHI, pH 8.0, 0.1% tween 20, 20 mM sodium EDTA) and rinsed with water. To achieve a dilute array, the beads were sonicated in 200 μl water and 2.5 μl evaporated onto a heated slide.

Enzyme Incorporation

A solution of the 51mer (SEQ ID NO. 3) (4 μM ; 2eqvs) in hybridisation buffer (5 mM, MgCl_2 , 7.5 mM DTT, 10 mM Tris.HCl (pH 7.6), 0.005% Triton X100) (20 ml) was added to 0.05 mg of beads (containing SEQ ID NO. 2) which were heated to 90°C for 2 min and allowed to cool for 1h. The fluorescent dUTP (400 μM stock, 0.5 μl , 10 μM ; 4eqvs) was added. A fraction of the beads were removed as a washing control and the polymerase (Sequence) (0.5 μl , 6.5 units) (one unit will incorporate 1 nmole dNTP in 30 s at 37°C) was added. The reaction was left at room temperature for 4 h and the beads were washed with NaOH, TT and TTE buffers as above and arrayed onto a coverslip.

The oligos used in this study are as follows:

3'-C(TAMRA)AGCGTCGGCAGGTATCCCAA-(C6amino)-5' SEQ ID NO. 1

and unlabelled:

5'-amino-GTCATCGAACGTCGAGCCTCGCAGCCGTCCAACCAACTCA-3'

SEQ ID NO. 2

10 and

3'-CAGTAGCTTGCAGCTCGGAGCGTCGGCAGGTTGGTTGAGTAGGTCTTGTTT-5'

SEQ ID NO. 3

15

as hybridised template.

Figure 1 shows the fluorescence image on the left and the optical image on the right when the experiment on the incorporation of fluorescently-labelled d-UTP was performed in the presence (A) and absence (B) of the polymerase. It is clear that no fluorescence is detected in the absence of any enzyme.

Figure 2 shows the beads diluted in unmodified beads so a quantitative analysis can be performed. The top figures (A) show the fluorescence from the beads with fluorophore-labelled DNA attached to the bead and the lower image (B) shows the level of fluorescence when the fluorophore-labelled nucleotide is incorporated into the DNA using a polymerase. The values of the fluorescence from the beads were compared:

(A) 3'-TAMRA DNA; Average counts/bead = 1956 (54 beads, +/- 50%)

30 (B) unblocked carboxylic acid-modified beads; counts/bead = 1739 (89%)
(88 beads, = +/- 40%)

This means the incorporation of the labelled nucleotide is 89%.

By comparing the signal-to-noise in Figure 1 between the level of fluorescence when the enzyme is present and when it is absent it is possible to estimate that the level of fluorescence could be reduced by a factor of 100-1000 while still allowing the detection of fluorescence above the background with adequate signal-to-noise. This



means that experiments can be performed with the fluorophore-labelled nucleotides highly diluted in non-labelled nucleotides so that only 1% of the fluorophore-labelled nucleotides are incorporated.

CLAIMS:

1. A method for determining the sequence of a target polynucleotide, comprising the steps of:
 - (i) contacting an array comprising multiple copies of the target polynucleotide with one of the bases A, T, G or C in a form and under conditions that permit polymerisation to occur, wherein a minor proportion of the base is detectably-labelled;
 - (ii) detecting the incorporation of the base onto the complement of the target after removal of non-incorporated base; and
 - (iii) repeating steps (i) and (ii) with each of the different bases until the sequence is determined.
2. A method according to claim 1, wherein the label is a fluorescent moiety.
3. A method according to claim 2, wherein step (ii) is carried out using optical means.
4. A method according to claim 3, wherein the optical means is a confocal scanning microscope.
5. A method according to any of claims 2 to 4, wherein step (ii) is carried out by measuring quantitatively the fluorescence signal generated on nucleotide incorporation.
6. A method according to any preceding claim, wherein the label is not removed from the incorporated nucleotides, and subsequent detection of incorporation is carried out by measuring the stepwise increase in the signal.
7. A method according to any of claims 1 to 5, wherein the label from incorporated bases is removed either prior to the addition of base having the same label or before it becomes difficult to detect incorporation.
8. A method according to claim 7, wherein the label is removed by photobleaching.
9. A method according to any preceding claim, wherein the proportion is less than 10%.
10. A method according to claim 9, wherein the proportion is less than 1%.
11. A method according to any preceding claim, wherein the polynucleotide is attached to the array via microspheres.
12. A method according to any preceding claim, wherein the detectably-labelled bases are dideoxynucleoside triphosphates.

Figure 1

(A)



(B)







Figure 2

(A)



(B)



SEQUENCE LISTING

<110> Solexa Ltd.

<120> POLYNUCLEOTIDE SEQUENCING

<130> REP06037WO

<140>

<141>

<150> 9907344.7

<151> 1999-03-30

<160> 3

<170> PatentIn Ver. 2.1

<210> 1

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<221> modified_base

<222> (1)..(21)

<223> n = cytosine with a fluorescent TMR group
attached.

m

= 5'-(C6amino)-adenine

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 1

nagcgtcggc aggtatccca m

21

<210> 2

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<221> modified_base

<222> (1)..(40)

<223> n = (amino)-guanine

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 2

ntcatcgaac gtcgagcctc gcagccgtcc aaccaactca

40

<210> 3

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 3

cagtagcttg cagctcggag cgtcggcagg ttggttgagt aggtcttggt t

51

INTERNATIONAL SEARCH REPORT

Int. Patent Application No.

PCT/GB 00/01222

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, CHEM ABS Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 90 13666 A (AMERSHAM INT PLC) 15 November 1990 (1990-11-15) the whole document ----	1-12
Y	HEAD ET AL.: "NESTED GENETIC BIT ANALYSIS (N-GBA) FOR MUTATION DETECTION IN THE p53 TUMOR SUPPRESSOR GENE" NUCLEIC ACIDS RESEARCH, vol. 25, no. 24, 1997, pages 5065-5071, XP002144794 the whole document ----	1-12
Y	WO 97 47761 A (SARNOFF CORP) 18 December 1997 (1997-12-18) the whole document ----- --/--	1-12



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

11 August 2000

Date of mailing of the international search report

24/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hagenmaier, S

INTERNATIONAL SEARCH REPORT

Int tional Application No
PCT/GB 00/01222

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 93 21340 A (MEDICAL RES COUNCIL ;BRENNER SYDNEY (GB); ROSENTHAL ANDRE (GB)) 28 October 1993 (1993-10-28) See page 12 line 6-line 35, page 14 line 1-line 29, page 19 line 31- page 20 line 4 the whole document ---	1-12
Y	US 5 547 839 A (FODOR STEPHEN P A ET AL) 20 August 1996 (1996-08-20) the whole document ---	1-12
A	US 5 302 509 A (CHEESEMAN PETER C) 12 April 1994 (1994-04-12) the whole document ---	
A	WO 91 06678 A (STANFORD RES INST INT ;TSIEN ROGER Y (US)) 16 May 1991 (1991-05-16) the whole document ---	
A	WO 93 05183 A (BAYLOR COLLEGE MEDICINE) 18 March 1993 (1993-03-18) the whole document ---	
A	WO 99 05321 A (TABONE JOHN C ;RAPIGENE INC (US); MOYNIHAN KRISTEN (US); NESS JEFF) 4 February 1999 (1999-02-04) the whole document ---	
A	EP 0 245 206 A (BATTELLE MEMORIAL INSTITUTE) 11 November 1987 (1987-11-11) the whole document ---	
E	WO 00 18956 A (SEEGER STEFAN) 6 April 2000 (2000-04-06) the whole document -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/01222

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9013666	A	15-11-1990	CA 2045505 A EP 0471732 A JP 4505251 T	12-11-1990 26-02-1992 17-09-1992
WO 9747761	A	18-12-1997	AU 3878497 A CA 2258511 A EP 0912752 A US 5908755 A	07-01-1998 18-12-1997 06-05-1999 01-06-1999
WO 9321340	A	28-10-1993	AT 159766 T AU 4020893 A CA 2133956 A DE 69314951 D DE 69314951 T EP 0640146 A ES 2110604 T JP 7507681 T US 6087095 A	15-11-1997 18-11-1993 28-10-1993 04-12-1997 19-03-1998 01-03-1995 16-02-1998 31-08-1995 11-07-2000
US 5547839	A	20-08-1996	US 5143854 A AU 9136791 A WO 9210587 A US 5902723 A AT 110738 T AT 175421 T AU 651795 B AU 5837190 A AU 672723 B AU 7765594 A BR 9007425 A CA 2054706 A DE 69012119 D DE 69012119 T DE 69032888 D DE 69032888 T DK 476014 T EP 0476014 A EP 0619321 A EP 0902034 A EP 0953835 A ES 2058921 T ES 2129101 T GB 2248840 A, B HK 61395 A HK 64195 A HU 59938 A IL 94551 A JP 11315095 A JP 11021293 A JP 4505763 T KR 9701577 B KR 9701578 B WO 9015070 A NL 191992 B NL 9022056 T NO 301233 B NZ 233886 A SG 13595 G	01-09-1992 08-07-1992 25-06-1992 11-05-1999 15-09-1994 15-01-1999 04-08-1994 07-01-1991 10-10-1996 04-05-1995 21-07-1992 08-12-1990 06-10-1994 22-12-1994 18-02-1999 29-07-1999 14-11-1994 25-03-1992 12-10-1994 17-03-1999 03-11-1999 01-11-1994 01-06-1999 22-04-1992 05-05-1995 05-05-1995 28-07-1992 30-03-1995 16-11-1999 26-01-1999 08-10-1992 11-02-1997 11-02-1997 13-12-1990 01-08-1996 02-03-1992 29-09-1997 25-02-1993 16-06-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/01222

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5547839 A		US 5744101 A	28-04-1998
		US 5489678 A	06-02-1996
		US 5889165 A	30-03-1999
		US 5753788 A	19-05-1998
		US 5744305 A	28-04-1998
		US 5770456 A	23-06-1998
		US 5800992 A	01-09-1998
		US 5424186 A	13-06-1995
		US 5405783 A	11-04-1995
		US 5871928 A	16-02-1999
US 5302509 A	12-04-1994	NONE	
WO 9106678 A	16-05-1991	CA 2044616 A	27-04-1991
		EP 0450060 A	09-10-1991
WO 9305183 A	18-03-1993	AU 2674092 A	05-04-1993
WO 9905321 A	04-02-1999	AU 8503298 A	16-02-1999
		EP 1000175 A	17-05-2000
EP 0245206 A	11-11-1987	AU 7583887 A	01-12-1987
		DK 688 A	17-02-1988
		WO 8706956 A	19-11-1987
		FI 875770 A	30-12-1987
		JP 1500221 T	26-01-1989
		NO 880010 A	10-02-1988
WO 0018956 A	06-04-2000	DE 19844931 C	15-06-2000
		AU 6197799 A	17-04-2000